

# Modern Approach to Lung Transplantation: Outcomes, Patient Selection & Waitlist management

**Nirmal S. Sharma M.D.**

*Medical Director, Lung Transplantation Program*

*Division of Pulmonary and Critical Care, Department of Medicine*

*Brigham and Women's Hospital & VA Boston Medical Center*

*Assistant Professor of Medicine, Harvard Medical School*

# Disclosures

- Scientific Advisory Board, Caredx
- Scientific Advisory Board, Resbiotic inc
- Scientific consultant, Sanofi
- Clinical Consultant/Medical Director, New England Donor Bank



Nirmal S Sharma MD

Medicine Residency @ Albert Einstein College of Medicine

Pulmonary & Critical Care Fellowship @ UAB Medicine

Lung transplantation & MCS training @ UAB Medicine

Assistant Professor of Medicine @HMS

Medical Director, Lung Transplantation @ BWH/VA Boston

- Clinical Focus: Lung transplantation and ECMO
- Research Focus: Immune mechanisms of allograft rejection

# Outline



- Epidemiology
  - Trends
  - Survival
- Organ Allocation System
- Recipient Selection
  - General Criteria
  - Disease specific considerations
- Transplant Waitlist Management
  - Donor Management
  - Candidate Management

# Epidemiology: Trends & Outcomes

# Epidemiology


- Trends
  - Increasing volume of transplants (mostly bilateral)
  - Increasing age of recipients
    - Mean age > 50 years
  - Increasing proportion of lung transplants for pulmonary fibrosis compared to COPD
  - Sicker candidates being transplanted

# Adult Lung Transplants by Year



OPTN/SRTR 2018 Annual Data Report: Lung. AJT

# Indications for Bilateral/Double Lung Transplants




**\*Other includes:**

Pulmonary Fibrosis, Other:	3.6%
Bronchiectasis:	4.0%
Sarcoidosis:	2.9%
Connective Tissue Disease:	1.5%
OB (non-Retx):	1.3%
LAM:	1.1%
Congenital Heart Disease:	1.1%
Miscellaneous:	2.0%

JHLT. 2014 Oct; 33(10): 1009-1024




# Adult Lung Transplants by Year



JHLT. 2018 Oct; 37(10): 1155-1206

# Adult Lung Transplants Survival


(Transplants: January 1990 – June 2016)




JHLT. 2018 Oct; 37(10): 1155-1206

# Adult CF Lung Transplants Survival

Survival Functions



Median Survival Time (Years)




Sethi J, Sharma NS. Ann Amer Thor Soc. 2020

# Overview of Allocation of Donor Lungs

# US organ allocation policy

- Pre-2005-
  - Based solely on amount of time on the waiting list
  - Often took several years to reach the top of the list
  - Led to strategy of time banking (premature listing to accrue time)
  - Biased against patients who were sicker or who had more rapidly progressive diseases
- Donor allocation by Lung Allocation Score (changed in 2005)
  - Geographic distribution
    - Allocation according to DSA, Zone A, B, C, D
    - Donor service area
      - Defined as area serviced by organ procurement organization
      - Brigham program gets priority access to donors from New England Organ Bank (NEOB)


# US organ allocation policy



- Change on Nov 27, 2017 to eliminate DSA prioritization
  - First level distribution now defined by a circle of radius 250 miles from donor hospital
  - Net migration of donors out of region
  - Increased patient acuity at time of transplantation

# Proposed continuous distribution of lungs

Consider multiple patient attributes all at once through a composite allocation score instead of within categories by sequence



# The Process of Lung Transplantation & Principals of Candidate Selection



# The Process of Lung Transplantation

Indications

Patient referral

Candidate selection

Waiting list

Donor organ selection

Transplant surgery

Peri-operative phase

Post-operative phase

Longterm follow-up

# Considerations for Lung Transplantation

- High (>50%) risk of death from lung disease within 2 years if lung transplant is not performed
- High (>80%) likelihood of surviving at least 90 days after transplant
- High (>80%) likelihood of 5-year post-transplant survival from a general medical health perspective, provided the graft functions properly

# Lung Transplant Referral: Practical Considerations

- NYHA class III or IV (ideally still ambulatory)
- Not severely debilitated
  - Absence of irreversible or untreatable extra-pulmonary disease
- Survival on waiting list depends on:
  - Waiting time (influenced by recipient size, blood group, presence of pre-formed antibodies)
  - Allocation method
  - Underlying disease

# Recipient Selection Criteria

- Untreatable end-stage parenchymal or pulmonary vascular disease
- No other significant medical diseases
- Substantial limitation of daily activity
- Limited life expectancy
- Ambulatory with rehabilitation potential
- Satisfactory nutritional, psychosocial profile and emotional support system

# Some Contraindications

- Significant extra-pulmonary organ dysfunction
- Malignancy (wait period up to 5 years after diagnosis based on stage and type of malignancy)
- Severe musculoskeletal disease
- Active substance addiction (including tobacco use)
- Untreatable psychosocial problems
- Medical non-compliance

# Lung Transplant Waitlist Management

# Challenges in Transplantation

- Donor shortage is the limiting factor in the number of transplants performed (20-25% of the offered lungs are accepted in the US)
- Inequities associated with donor allocation system
- Average waiting time varies by region
- As disease progresses, physical conditioning worsens often making a candidate too sick for transplant.

# Increasing Organ Availability




# Extended Donor Guidelines

- Age < 60-65 years
- No significant lung disease
- Limited smoking history (< 20 pk-years)
- Clear chest radiograph
- Gas exchange  $\text{PaO}_2/\text{FiO}_2 > 300$  on 5 cm  $\text{H}_2\text{O}$  PEEP
- Absence of secretions / aspiration
- Normal endobronchial exam by bronchoscopy
- Exclusion: active infection, HIV, hepatitis, malignancy

# Optimizing Donor Lungs

**Table 1** Lung Donor-Management Protocol

1. Apnea test performed with ventilator (continuous positive pressure mode)
2. Mechanical ventilation with positive end-expiratory pressure 8 to 10 cm H<sub>2</sub>O and tidal volume of 6–8 ml/kg
3. Recruitment maneuvers once per hour and after any disconnection from the ventilator.
4. Bronchoscopy with bilateral bronchoalveolar lavage.
5. Hemodynamics closely monitored with PICCO System, goal of extravascular lung water < 10 ml/kg (administering diuretics if necessary), and central venous pressure objective < 8 mm Hg
6. Methylprednisolone (15 mg/kg) after brain death declaration
7. Alveolar recruitment involved controlled ventilation (peak pressure limit of 35 mm Hg) with positive end-expiratory pressure of 18–20 cm H<sub>2</sub>O for 1 minute, and decreased 2 cm H<sub>2</sub>O each minute; after that we increased 50% tidal volumes for 10 breaths




**Figure 1** Lung donation effectiveness at University Hospital Marqués de Valdecilla, 2003 to 2011.

JHLT 2014; 33: 178


# Increasing Donor Lung Availability

## Donation after cardiac death



**2001 – 2010: Overall increase of 500%**

## Donation after brain death




**2001 – 2010: Overall decrease of 25%**

*Courtesy of New England Organ Bank*

# Ex vivo Lung Perfusion

## Ex vivo Lung Perfusion Equipment



**Table 3**  
Ventilatory and perfusion strategy for EVLP

### Ventilation

Tidal volume	7 mL/kg
PEEP	5 cm H <sub>2</sub> O
Frequency	7 breaths/min
I/E ratio	1/2
Recruitment	1 every hour to PawP 20 cm H <sub>2</sub> O

### Perfusion

Pump flow	40% estimated donor cardiac output
Pulmonary artery pressure	7–13 mm Hg <sup>a</sup>
Left atrial pressure	3–5 mm Hg <sup>b</sup>
Perfusate exchange	250 mL every hour
Perfusate composition	Steen solution, heparin, antibiotics, solumedrol
Perfusate pH	6.8–7.4
Perfusate Pco <sub>2</sub>	35–45 mm Hg

Clin Chest Med 2011; 32: 233-44

ORIGINAL ARTICLE

# Heart and Lung Transplants from HCV-Infected Donors to Uninfected Recipients

Ann E. Woolley, M.D., Steve K. Singh, M.D., Hilary J. Goldberg, M.D., Hari R. Mallidi, M.D., Michael M. Givertz, M.D., Mandeep R. Mehra, M.D., Antonio Coppolino, M.D., Amanda E. Kuzstos, B.S., Megan E. Johnson, B.A., Kaiwen Chen, B.S., Esther A. Haddad, M.D., John Fanikos, R.Ph., David P. Harrington, Ph.D., Phillip C. Camp, M.D., and Lindsey R. Baden, M.D., for the DONATE HCV Trial Team\*






Figure 2. Results of Patient Follow-up after Transplantation.

The black X indicates that Patient 1 died at week 32. The green bars represent patients who had not completed 16 weeks of follow-up by July 31, 2018.

# Candidate Management

# Optimizing Composite Allocation Score

- <http://optn.transplant.hrsa.gov/resources/all-oculationcalculators.asp?index=88>


Date of birth: *	<input type="text"/> 	Date added to the waiting list: *	<input type="text"/> 
Height: *	<input type="text"/> ft <input type="text"/> in <input type="text"/> cm	Weight: *	<input type="text"/> lbs <input type="text"/> kg
Blood type: *	<input type="text"/>	Prior living donor: *	<input type="radio"/> Yes <input type="radio"/> No
Lung diagnosis code *	<input type="text"/>	Detailed CPRA: *	<input type="text"/> <a href="#">CPRA Calculator</a>
Functional status:	<input type="text"/>	Assisted ventilation:	<input type="text"/>
Requires supplemental O <sub>2</sub> :	<input type="text"/>	Pulmonary artery systolic pressure:	<input type="text"/> mmHg
Mean pulmonary artery pressure:	<input type="text"/> mmHg	Cardiac index (CI):	<input type="text"/> L/min/m <sup>2</sup>
6-minute walk distance:	<input type="text"/> feet	Total bilirubin:	<input type="text"/> mg/dL
Serum creatinine:	<input type="text"/> mg/dL		
<p><b>i</b> If using a central venous test value for PCO<sub>2</sub> subtract 6 mmHg before entering the value.</p>			
PCO <sub>2</sub> :			
Current:			
<input type="text"/> mmHg			
Highest:			
<input type="text"/> mmHg			
Lowest:			
<input type="text"/> mmHg			

# ECMO as a Bridge to Transplant

J Thorac Cardiovasc Surg. 2013 Mar;145(3):862-7; discussion 867-8. doi: 10.1016/j.jtcvs.2012.12.022. Epub 2013 Jan 11.


## Extracorporeal membrane oxygenation as a bridge to pulmonary transplantation.

Hoopes CW<sup>1</sup>, Kukreja J, Golden J, Davenport DL, Diaz-Guzman E, Zwischenberger JB.



- Average duration of extracorporeal membrane oxygenation support was 13.7 days (range, 2-53 days),
- Mean survival of all patients bridged to pulmonary transplantation was 26 months (range, 54 days to 95 months).
- The 1-, 3-, and 5-year survival was 93%, 80%, and 66%





Thor CV Surg 2013; 145: 862









# Summary

- Lung transplantations are increasing yearly
- Changes in the organ allocations system are focused on balancing transplantation of sick recipients but also those who have long term survivability.
- Optimal management requires a multidisciplinary approach
- Multimodal approach used to augment organ availability and reduce waitlist mortality.

# Thank You

## Questions?

